



## TH gene

tyrosine hydroxylase

### Normal Function

The *TH* gene provides instructions for making the enzyme tyrosine hydroxylase, which is important for normal functioning of the nervous system. Tyrosine hydroxylase takes part in the first step of the pathway that produces a group of hormones called catecholamines. This enzyme helps convert the protein building block (amino acid) tyrosine to a catecholamine called dopamine. Dopamine is also known as a neurotransmitter because it transmits signals between nerve cells in the brain to help control physical movement and emotional behavior. Other catecholamines called norepinephrine and epinephrine are produced from dopamine. Norepinephrine and epinephrine are involved in the autonomic nervous system, which controls involuntary body processes such as the regulation of blood pressure and body temperature.

### Health Conditions Related to Genetic Changes

#### dopa-responsive dystonia

More than two dozen mutations in the *TH* gene have been found to cause dopa-responsive dystonia. This condition is characterized by a pattern of involuntary muscle contractions (dystonia), tremors, and other uncontrolled movements and usually responds to treatment with a medication called L-Dopa. Most *TH* gene mutations that cause this condition change single protein building blocks (amino acids) in the tyrosine hydroxylase enzyme, resulting in a decrease in functional enzyme. A reduction in normal tyrosine hydroxylase enzyme leads to a decrease in the production of dopamine, which causes the movement problems characteristic of dopa-responsive dystonia. The amount of functional enzyme that is produced is associated with the severity of the signs and symptoms. Less functional enzyme leads to more severe symptoms.

#### tyrosine hydroxylase deficiency

More than 20 mutations in the *TH* gene have been identified in people with tyrosine hydroxylase (TH) deficiency. These mutations result in reduced activity of the tyrosine hydroxylase enzyme. As a result, the body produces less dopamine, norepinephrine, and epinephrine. These catecholamines are necessary for normal nervous system function, and changes in their levels contribute to the abnormal movements, nervous system dysfunction, and other neurological problems seen in people with TH deficiency.

Dopa-responsive dystonia is sometimes considered a mild form of tyrosine hydroxylase deficiency. It is uncertain whether they are two separate disorders or part of the same disease spectrum.

### other disorders


Certain common *TH* variations (polymorphisms) modify catecholamine production, which affects the risk of developing conditions associated with regulation of the autonomic nervous system. These *TH* gene polymorphisms affect the extent to which blood pressure increases with stress and may increase the risk of high blood pressure (hypertension).

One *TH* gene polymorphism has been associated with sudden infant death syndrome (SIDS). SIDS is a major cause of death in babies younger than 1 year. It is characterized by sudden and unexplained death, usually during sleep. The polymorphism, called allele \*9.3, is the most common *TH* gene polymorphism among people of European descent and has been identified in a larger percentage of babies who die from SIDS than in other babies. This version of the gene may affect the regulation of breathing or awakening in infants.

### **Chromosomal Location**

Cytogenetic Location: 11p15.5, which is the short (p) arm of chromosome 11 at position 15.5

Molecular Location: base pairs 2,163,929 to 2,174,081 on chromosome 11 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

### **Other Names for This Gene**

- DYT5b
- TY3H\_HUMAN
- TYH
- tyrosine 3-monooxygenase

## **Additional Information & Resources**

### Educational Resources

- Basic Neurochemistry (6th edition, 1999): Tyrosine hydroxylase is the rate-limiting enzyme for the biosynthesis of catecholamines  
<https://www.ncbi.nlm.nih.gov/books/NBK27988/#A861>
- Neuroscience (second edition, 2001): Regulation of tyrosine hydroxylase by protein phosphorylation (image)  
<https://www.ncbi.nlm.nih.gov/books/NBK10990/figure/A576/>

### GeneReviews

- Tyrosine Hydroxylase Deficiency  
<https://www.ncbi.nlm.nih.gov/books/NBK1437>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28tyrosine+hydroxylase%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D>

### OMIM

- TYROSINE HYDROXYLASE  
<http://omim.org/entry/191290>

### Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC\\_TH.html](http://atlasgeneticsoncology.org/Genes/GC_TH.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=TH%5Bgene%5D>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=11782](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=11782)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/7054>
- UniProt  
<http://www.uniprot.org/uniprot/P07101>

## Sources for This Summary

- Furukawa Y, Kish SJ, Fahn S. Dopa-responsive dystonia due to mild tyrosine hydroxylase deficiency. *Ann Neurol*. 2004 Jan;55(1):147-8.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/14705130>
- GeneReview: Tyrosine Hydroxylase Deficiency  
<https://www.ncbi.nlm.nih.gov/books/NBK1437>
- Hoffmann GF, Assmann B, Bräutigam C, Dionisi-Vici C, Häussler M, de Klerk JB, Naumann M, Steenbergen-Spanjers GC, Strassburg HM, Wevers RA. Tyrosine hydroxylase deficiency causes progressive encephalopathy and dopa-nonresponsive dystonia. *Ann Neurol*. 2003;54 Suppl 6: S56-65.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12891655>
- Hunt SC. Tyrosine hydroxylase: another piece of the genetics of hypertension puzzle. *Circulation*. 2007 Aug 28;116(9):970-2.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/17724269>
- Klitschar M, Reichenpfader B, Saternus KS. A functional polymorphism in the tyrosine hydroxylase gene indicates a role of noradrenalinergic signaling in sudden infant death syndrome. *J Pediatr*. 2008 Aug;153(2):190-3. doi: 10.1016/j.jpeds.2008.02.032. Epub 2008 Apr 3.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18534229>
- Pearl PL, Capp PK, Novotny EJ, Gibson KM. Inherited disorders of neurotransmitters in children and adults. *Clin Biochem*. 2005 Dec;38(12):1051-8. Epub 2005 Nov 18. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16298354>
- Pearl PL, Taylor JL, Trzcinski S, Sokohl A. The pediatric neurotransmitter disorders. *J Child Neurol*. 2007 May;22(5):606-16. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/17690069>
- Rao F, Zhang L, Wessel J, Zhang K, Wen G, Kennedy BP, Rana BK, Das M, Rodriguez-Flores JL, Smith DW, Cadman PE, Salem RM, Mahata SK, Schork NJ, Taupenot L, Ziegler MG, O'Connor DT. Adrenergic polymorphism and the human stress response. *Ann N Y Acad Sci*. 2008 Dec;1148: 282-96. doi: 10.1196/annals.1410.085.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/19120120>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2743085/>
- Rao F, Zhang L, Wessel J, Zhang K, Wen G, Kennedy BP, Rana BK, Das M, Rodriguez-Flores JL, Smith DW, Cadman PE, Salem RM, Mahata SK, Schork NJ, Taupenot L, Ziegler MG, O'Connor DT. Tyrosine hydroxylase, the rate-limiting enzyme in catecholamine biosynthesis: discovery of common human genetic variants governing transcription, autonomic activity, and blood pressure in vivo. *Circulation*. 2007 Aug 28;116(9):993-1006. Epub 2007 Aug 13.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/17698732>
- Schiller A, Wevers RA, Steenbergen GC, Blau N, Jung HH. Long-term course of L-dopa-responsive dystonia caused by tyrosine hydroxylase deficiency. *Neurology*. 2004 Oct 26;63(8):1524-6.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15505183>
- OMIM: TYROSINE HYDROXYLASE  
<http://omim.org/entry/191290>

- Verbeek MM, Steenbergen-Spanjers GC, Willemsen MA, Hol FA, Smeitink J, Seeger J, Grattan-Smith P, Ryan MM, Hoffmann GF, Donati MA, Blau N, Wevers RA. Mutations in the cyclic adenosine monophosphate response element of the tyrosine hydroxylase gene. *Ann Neurol*. 2007 Oct;62(4):422-6.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/17696123>
- Yeung WL, Wong VC, Chan KY, Hui J, Fung CW, Yau E, Ko CH, Lam CW, Mak CM, Siu S, Low L. Expanding phenotype and clinical analysis of tyrosine hydroxylase deficiency. *J Child Neurol*. 2011 Feb;26(2):179-87. doi: 10.1177/0883073810377014. Epub 2010 Sep 7. Erratum in: *J Child Neurol*. 2012 Jun;27(6):829-31.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/20823027>
- Zhang L, Rao F, Wessel J, Kennedy BP, Rana BK, Taupenot L, Lillie EO, Cockburn M, Schork NJ, Ziegler MG, O'Connor DT. Functional allelic heterogeneity and pleiotropy of a repeat polymorphism in tyrosine hydroxylase: prediction of catecholamines and response to stress in twins. *Physiol Genomics*. 2004 Nov 17;19(3):277-91. Epub 2004 Sep 14.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15367723>

---

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/TH>

Reviewed: May 2012  
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services